

105 學年 第一學期

離散數學 第二次平時考試 資訊二乙

學號：

姓名：Solutions

1. (20 points) (a) 集合  $S$  的定義如下，寫出  $S$  的幕集合 (power set)  $\mathcal{P}(S)$ 。(Set  $S$  is defined as below. Write the power set of  $S$ ,  $\mathcal{P}(S)$ .)

$$S = \{x, y, \{x, y\}\}$$

- (b) 假設  $A, B,$  和  $C$  是三個  $\mathcal{P}(S)$  中的不同元素 (elements)，且  $A \cup (B \cap C) = \{x, y\}$ 。寫出三組不同的  $A, B,$  和  $C$ 。(Let  $A, B,$  and  $C$  be three various elements in  $\mathcal{P}(S)$  and  $A \cup (B \cap C) = \{x, y\}$ . Write three different solutions of  $A, B,$  and  $C$ .)

(a)  $\mathcal{P}(A) = \{\emptyset, \{x\}, \{y\}, \{\{x, y\}\}, \{x, y\}, \{x, \{x, y\}\}, \{y, \{x, y\}\}, \{x, y, \{x, y\}\}$ .

(b) (i)  $A = \{x\}, B = \{y\}, C = \{x, y\}$ ; (ii)  $A = \{x, y\}, B = \{x\}, C = \{y\}$ ; (iii)  $A = \emptyset, B = \{x, y\}, C = \{x, y, \{x, y\}\}$ .

2. (30 points) 假設  $g$  是自  $A$  映到  $B$  的函數及  $f$  是  $B$  映到  $C$  的函數，分別考慮下列兩個敘述。如果該敘述為真，則證明之；否則，則舉出一個反例。(Let  $g$  be a function from  $A$  to  $B$  and  $f$  be a function  $B$  to  $C$ . For each of the following two statements, if it is true, prove; otherwise, give a counterexample.)

- (a) 若  $f \circ g$  是一個映成函數，則  $f$  和  $g$  都是映成函數。(if  $f \circ g$  is an onto function,  $f$  and  $g$  are onto functions.)

- (b) 若  $f$  和  $g$  都是一對一函數，則  $f \circ g$  是一個一對一函數。(if  $f$  and  $g$  are one-to-one functions,  $f \circ g$  is a one-to-one function.)

(a) Let  $A = \{1, 2, 3\}, B = \{a, b, c, d\}, C = \{x, y, z\}$ . If  $g(1) = a, g(2) = b,$  and  $g(3) = c,$  then  $g$  is not an onto function; if  $f(a) = x, f(b) = y, f(c) = z,$  and  $f(d) = z,$  then  $f$  is an onto function. However,  $f \circ g(1) = x, f \circ g(2) = y,$  and  $f \circ g(3) = z$  such that  $f \circ g$  is an onto function. We have a counterexample that  $f \circ g$  is an onto function, but  $g$  is not.

(b) Let  $a$  and  $b$  be two distinct elements of  $A$  and  $g(a) = r$  and  $g(b) = s$ . Since  $g$  is a one-to-one function,  $r$  and  $s$  must be two distinct elements of  $B$ . Let  $f(r) = x$  and  $f(s) = y$ . Also, since  $f$  is a one-to-one function,  $x$  and  $y$  must be two distinct elements of  $C$ . For distinct elements  $a$  and  $b$  of  $A$ , We have  $f \circ g(a) = x, f \circ g(b) = y,$  and  $x \neq y$ . Hence,  $f \circ g$  is a one-to-one function from  $A$  to  $C$ .

3. (20 points) 假設  $A$  和  $B$  是兩個集合，證明  $(A \cup B) \cap (A \cup \bar{B}) = A$ 。(Let  $A$  and  $B$  be two sets, prove  $(A \cup B) \cap (A \cup \bar{B}) = A$ .)

$$A = A \cup \emptyset$$

(Identity Law)

$$= A \cup (B \cap \bar{B})$$

(Complement Law)

$$= (A \cup B) \cap (A \cup \bar{B})$$

(Distributive Law)

4. (30 points) (a) 定義對等關係 (equivalence relation) (Define equivalence relation.)

- (b) 假設  $f: A \rightarrow B$  是一個函數，關係  $R = \{(x, y) \mid x, y \in A \text{ 且 } f(x) = f(y)\}$ ；證明  $R$  是一個對等關係。(Let  $f: A \rightarrow B$  be a function and relation  $R = \{(x, y) \mid x, y \in A \text{ and } f(x) = f(y)\}$ . Prove  $R$  is an equivalence relation.)

- (c) 如果 (b) 小題的  $f$  是一個整數到整數的函數，且  $f(x) = x^2$ ，寫出關係  $R$  所產生的對等類別  $[a]_R, a \in A$ 。(Suppose  $f$  in (b) is a function from  $Z$  to  $Z$  and  $f(x) = x^2$ . Write the equivalence class  $[a]_R$ , generated by relation  $R, a \in A$ .)

- (a) A relation  $R \subseteq A \times A$  is an equivalence relation if it satisfies the following laws:

i. Reflexive law:  $x \in A \Rightarrow (x, x) \in R$ .

ii. Symmetric law:  $x, y \in A \wedge (x, y) \in R \Rightarrow (y, x) \in R$ .

iii. Transitive law:  $x, y, z \in A \wedge (x, y) \in R \wedge (y, z) \in R \Rightarrow (x, z) \in R$ .

- (b) Show  $R$  is and equivalence equation.

- i. Reflexive law: For an element  $x$  of  $A$ , since  $f(x)=f(x)$ , we have  $(x, x) \in R$ .
  - ii. Symmetric law: For  $x, y \in A \wedge$  if  $(x, y) \in R$ , then we have  $f(x)=f(y)$ . Hence  $f(y)=f(x)$ , we obtain  $(y, x) \in R$ .
  - iii. Transitive law: For  $x, y, z \in A \wedge (x, y) \in R \wedge (y, z) \in R$ , i.e.,  $f(x)=f(y)$  and  $f(y)=f(z)$ . Thus, we have  $f(x)=f(z)$  and conclude  $(x, z) \in R$ .
- (c)  $[a]_R = \{x \mid x \in A \wedge |x|=a\}$ , e.g.,  $[0]_R = \{0\}$ ,  $[1]_R = \{-1, 1\}$ ,  $[n]_R = \{-n, n\}$ .